

# REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank) 2. REPORT DATE 2/21/96 3. REPORT TYPE AND DATES COVERED Final Report 12/16/94-12/14/95

## 4. TITLE AND SUBTITLE

Study of Anion-Molecule Reaction Dynamics with Time-Dependent Photoelectron Spectroscopy

## 5. FUNDING NUMBERS

61103D  
3484/05

## 6. AUTHOR(S)

Daniel M. Neumark

AFOSR-TR-96

## 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)

University of California  
Sponsored Projects Office  
336 Sproul Hall  
Berkeley, California 94720-5940

0122

## 9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)

Air Force Office of Scientific Research/NL  
Chemistry and Materials Science Research  
110 Duncan Avenue, Suite B115  
Bolling Air Force Base, DC 20332-0001

## 10. SPONSORING / MONITORING AGENCY REPORT NUMBER

F49620-95-1-0078

## 11. SUPPLEMENTARY NOTES

## 12a. DISTRIBUTION / AVAILABILITY STATEMENT

Approved for public release; distribution is unlimited

## 12b. DISTRIBUTION CODE

## 13. ABSTRACT (Maximum 200 words)

A novel femtosecond time-resolved experiment has been set up to probe the photodissociation dynamics of negative ions and negative ion clusters. First results for the photodissociation of  $I_2^-$  have been obtained recently.

19960320 041

## 14. SUBJECT TERMS

negative ions, photodissociation, femtosecond photoelectron spectroscopy

## 15. NUMBER OF PAGES

3

## 16. PRICE CODE

## 17. SECURITY CLASSIFICATION OF REPORT

UNCLASSIFIED

## 18. SECURITY CLASSIFICATION OF THIS PAGE

UNCLASSIFIED

## 19. SECURITY CLASSIFICATION OF ABSTRACT

UNCLASSIFIED

## 20. LIMITATION OF ABSTRACT

UNLIMITED (SAR)

## FINAL TECHNICAL REPORT

**TITLE:** Study of Anion-Molecule Reaction Dynamics with Time-Dependent Photoelectron Spectroscopy

**PRINCIPAL INVESTIGATOR:** Daniel M. Neumark

**DATE:** December 16, 1994 - December 14, 1995

**GRANT NO:** F4960-95-1-0078


### Abstract:

A novel femtosecond time-resolved experiment has been set up to probe the photodissociation dynamics of negative ions and negative ion clusters. First results for the photodissociation of  $I_2^-$  have been obtained recently.

### Final Technical Report:

DURIP funds were used to purchase a Ti:sapphire femtosecond laser system (Model #CPA-1000 MPS) from Clark MRX Inc., at a cost of \$165,600. The Ti:sapphire femtosecond laser system is being used on a new experiment designed to study the photodissociation dynamics of negative ions and negative ion clusters on a femtosecond time scale. The novel feature of this experiment is that photoelectron spectroscopy is used to probe the dynamics of the dissociating anion in real-time. We have very recently obtained our first results on  $I_2^-$  photodissociation.

The principle of the experiment can be best understood by specifically considering its application to  $I_2^-$ :



This is a pump-probe experiment using two femtosecond laser pulses. The pump pulse ( $h\nu_1$ , 780 nm) excites mass-selected  $I_2^-$  from the ground  $X^2\Sigma_u^+$  state to the repulsive  $2\Pi_{u(3/2)}$  state. The  $I_2^-$  begins to dissociate, and after a variable time delay  $\Delta t$ , the dissociating ions are photodetached with the probe pulse ( $h\nu_2$ , 260 nm). We then measure the photoelectron spectrum, thereby mapping out the dissociating wave packet onto the well-known  $I_2$  potential energy surfaces.

At short times, one expects to see a transient associated with the dissociating  $I_2^-$  molecule, and at longer times, the  $I^-$  photoelectron spectrum should become more prominent as dissociation occurs. This is precisely what has been observed in our first measurements on  $I_2^-$ . However, the time scale for dissociation appears to be several hundred fsec longer than predicted by our simulations that use the currently accepted potential energy curves for  $I_2^-$ , indicating that these potentials may require significant modification. This issue will be addressed in further experiments and theoretical work.